


RESEARCH ARTICLE

Open Access



# Ca<sup>2+</sup> mobilization-dependent reduction of the endoplasmic reticulum lumen is due to influx of cytosolic glutathione

Beáta Lizák<sup>1\*</sup> , Julia Birk<sup>2</sup>, Melinda Zana<sup>3,4</sup>, Gergely Kosztyi<sup>1</sup>, Denise V. Kratschmar<sup>2</sup>, Alex Odermatt<sup>2</sup>, Richard Zimmermann<sup>5</sup>, Miklós Geiszt<sup>3,4</sup>, Christian Appenzeller-Herzog<sup>2,6\*</sup> and Gábor Bánhegyi<sup>1</sup>

## Abstract

**Background:** The lumen of the endoplasmic reticulum (ER) acts as a cellular Ca<sup>2+</sup> store and a site for oxidative protein folding, which is controlled by the reduced glutathione (GSH) and glutathione-disulfide (GSSG) redox pair. Although depletion of luminal Ca<sup>2+</sup> from the ER provokes a rapid and reversible shift towards a more reducing poise in the ER, the underlying molecular basis remains unclear.

**Results:** We found that Ca<sup>2+</sup> mobilization-dependent ER luminal reduction was sensitive to inhibition of GSH synthesis or dilution of cytosolic GSH by selective permeabilization of the plasma membrane. A glutathione-centered mechanism was further indicated by increased ER luminal glutathione levels in response to Ca<sup>2+</sup> efflux. Inducible reduction of the ER lumen by GSH flux was independent of the Ca<sup>2+</sup>-binding chaperone calreticulin, which has previously been implicated in this process. However, opening the translocon channel by puromycin or addition of cyclosporine A mimicked the GSH-related effect of Ca<sup>2+</sup> mobilization. While the action of puromycin was ascribable to Ca<sup>2+</sup> leakage from the ER, the mechanism of cyclosporine A-induced GSH flux was independent of calcineurin and cyclophilins A and B and remained unclear.

**Conclusions:** Our data strongly suggest that ER influx of cytosolic GSH, rather than inhibition of local oxidoreductases, is responsible for the reductive shift upon Ca<sup>2+</sup> mobilization. We postulate the existence of a Ca<sup>2+</sup>- and cyclosporine A-sensitive GSH transporter in the ER membrane. These findings have important implications for ER redox homeostasis under normal physiology and ER stress.

**Keywords:** Endoplasmic reticulum, Endoplasmic reticulum stress, Redox homeostasis, Glutathione, Calcium, Sec61 translocon, Cyclosporine A, Cyclophilins, Calreticulin, Membrane transport proteins

## Background

The lumen of the endoplasmic reticulum (ER) is the first compartment of the eukaryotic secretory pathway. Its content resembles that of an “extracellular space inside the cell.” For example, it is characterized by a high Ca<sup>2+</sup> concentration and an oxidizing redox balance [1–3],

whereas the term “redox balance” shall herein refer to the thiol/disulfide system only.

Proper maintenance of the intraluminal homeostasis in the ER is a vital requirement for the cell. Either the depletion of luminal Ca<sup>2+</sup> or the alteration of the redox balance can lead to ER stress that is an ominous accumulation of unfolded proteins in the ER lumen. ER stress triggers an adaptive program of signal transduction pathways, called the Unfolded Protein Response (UPR). Unresolved ER stress can finally result in programmed cell death [4].

The ER lumen serves as the main source of releasable Ca<sup>2+</sup> for cytosolic signaling, which is maintained by the Sarcoplasmic/Endoplasmic Reticulum Calcium ATP-ase

\* Correspondence: [lizak.beata@med.semmelweis-univ.hu](mailto:lizak.beata@med.semmelweis-univ.hu); [christian.appenzeller@unibas.ch](mailto:christian.appenzeller@unibas.ch)

<sup>1</sup>Department of Medical Chemistry, Molecular Biology and Pathobiochemistry, Semmelweis University, Budapest, Hungary

<sup>2</sup>Division of Molecular and Systems Toxicology, Department of Pharmaceutical Sciences, University of Basel, Klingelbergstrasse 50, 4056 Basel, Switzerland

Full list of author information is available at the end of the article



(SERCA) pump. SERCA-dependent  $\text{Ca}^{2+}$  influx is counter-balanced by a basal  $\text{Ca}^{2+}$  leakage and the opening of various second messenger gated channels activated by different extracellular stimuli [5]. Besides being the  $\text{Ca}^{2+}$  store, the high luminal  $\text{Ca}^{2+}$  concentration is indispensable for the function of critical components of the protein folding machinery such as chaperones and folding enzymes [6, 7].

Formation of native disulfide bonds in secretory and membrane proteins is a crucial step in protein maturation. Oxidation of cysteine residues in nascent polypeptides or rearrangement of misplaced disulfide bonds is catalyzed by the members of the Protein Disulfide Isomerase (PDI) family, the reoxidation of which can happen through various pathways [8]. The reduced glutathione (GSH) and glutathione-disulfide (GSSG) redox pair is the major low molecular weight thiol-disulfide buffer in the ER lumen [9]. Both GSH and GSSG were shown to directly react with the active centers of PDIs [10]. Total glutathione concentration in the ER reaches millimolar ranges providing an exceptionally high buffering capacity against oxidizing or reducing imbalances [9, 11, 12].

Participating in second-order thiol-disulfide exchange reactions, the reducing power of glutathione depends on  $[\text{GSH}]^2:[\text{GSSG}]$  rather than on the bimolecular ratio  $[\text{GSH}]:[\text{GSSG}]$  [9, 10]. The  $[\text{GSH}]^2:[\text{GSSG}]$  ratio in the ER lumen is far more oxidizing than the cytosolic redox poise [13, 14]. This is also reflected by a higher  $[\text{GSH}]:[\text{GSSG}]$  [15]. The most recent estimation of ER luminal  $[\text{GSH}]:[\text{GSSG}]$  derives from intact HeLa cells using the glutathionylation state of a single cysteine mutant glutaredoxin, which calculated a bimolecular ratio of less than 7:1 [12]. According to these numbers, the ER luminal glutathione concentration  $[\text{GSH}] + 2[\text{GSSG}]$  is twofold higher than the total cellular glutathione concentration [12].

The source of ER luminal GSH has to be the cytosolic glutathione pool, because the ER is devoid of enzymes for GSH synthesis. GSH was indeed shown to permeate the ER membrane; a facilitated diffusion selective to GSH was described in rat liver microsomes. On the other hand, microsomes were impermeable for GSSG, which was entrapped in the lumen upon GSH addition [16]. GSH permeation from the cytosol was also confirmed by showing the direct modification of luminal oxidoreductases by GSH [17, 18]. GSH can be directly oxidized by many intraluminal reactions involving the oxidative protein folding machinery; thus, the  $[\text{GSH}]:[\text{GSSG}]$  ratio is constantly shifted towards the oxidized form. The locally accumulated GSSG can leave the ER through the secretory pathway or can also react with reduced PDI for subsequent disulfide bond formation in client proteins [19, 20].

Buffering the luminal  $[\text{GSH}]^2:[\text{GSSG}]$  ratio is indispensable for correct disulfide bond formation; therefore, it is strictly regulated by luminal oxidoreductases [21].

An over-oxidizing environment can lead to unwanted disulfide bond formation, which in turn can provoke the UPR or, in serious cases, apoptosis [22]. On the contrary, an over-reducing environment prevents disulfide bond formation and protein secretion; however, it can help the clearance of misfolded polypeptides. Since the maintenance of a proper redox distribution in PDIs active sites depends on the reducing power of GSH [17], the control of GSH uptake from the cytosol can be an important question.

Recently, several groups reported a reducing shift of the ER luminal redox balance upon  $\text{Ca}^{2+}$  depletion [13, 23, 24]. Inhibition of  $\text{Ca}^{2+}$  uptake by SERCA pump or hormones inducing ER  $\text{Ca}^{2+}$  release caused immediate reduction of the ER lumen. Biophysically different redox-sensitive fluorescent readouts like the fluorescent lifetime of roGFPiE [23] or the excitation ratio of Grx1-roGFP1-iE<sub>ER</sub> [25] or an OxyR-YFP fusion protein called HyPer-ER [24] observed the same phenomenon in living cell experiments. Whereas roGFPiE reacts with thiol-disulfide couples [23] and HyPer-ER with thiol-disulfide couples or  $\text{H}_2\text{O}_2$  with unclear specificities [26, 27], Grx1-roGFP1-iE<sub>ER</sub> is a bona fide  $[\text{GSH}]^2:[\text{GSSG}]$  sensor [13]. A direct  $\text{Ca}^{2+}$  sensitivity of the probes was also excluded [23], suggesting that indeed  $\text{Ca}^{2+}$  cues can physiologically regulate the ER redox balance. The rapid reductive shift can be explained by a quick change of the local concentrations of redox active compounds, either by the uptake of reducing or by the release of oxidizing molecules. Furthermore, rapid activation/inhibition of ER oxidoreductases upon  $\text{Ca}^{2+}$  depletion might also influence the luminal redox balance. In this vein, it was supposed that selective sequestration of PDI1A with calreticulin (CRT) in a complex which formed under  $\text{Ca}^{2+}$ -depleted conditions decreases the effective concentration of this major thiol oxidant, resulting in a hypo-oxidizing state [28]. The same study also showed that the major ER thiol oxidase ERO1 was insensitive to changes in  $[\text{Ca}^{2+}]$  [28].

$\text{Ca}^{2+}$  mobilization also triggers a rapid increase in [ATP] in the ER lumen [29]. The underlying mechanism involves the ER membrane ATP/ADP exchanger AXER, which increases ATP import following enhanced glycolytic flux downstream of a  $\text{Ca}^{2+}$ -dependent CAMKK2-AMPK signaling cascade [30], and a likely temporary lowering of ATP consumption in the ER in response to ER  $\text{Ca}^{2+}$  depletion [31]. Still, the molecular identification of most of the transporter proteins in the ER membrane is still missing, although biochemical evidence describing many carrier-mediated transport processes is available [32]. Non-specific membrane permeation possibilities also exist, for example, the translocon polypeptide channel was described as a pore in the ER membrane allowing the transition of ions including  $\text{Ca}^{2+}$ , and several

small molecules [33, 34]. The permeability of the translocon pore is known to be regulated by BiP, the most prominent chaperone of the ER lumen being a  $\text{Ca}^{2+}$ -binding protein itself [35]. A recent study reported that the translocon in yeast can mediate GSH influx into the ER and that the channel is gated by oxidized Kar2, the yeast orthologue of BiP [36].

In this study, we further examined the mechanism of  $\text{Ca}^{2+}$ -sensitive reduction of the ER lumen by real-time measurements using Grx1-roGFP1-iE<sub>ER</sub> and HyPer-ER and found evidence for a  $\text{Ca}^{2+}$  depletion-driven GSH transport process through the ER membrane.

## Results

### Reduction of the ER lumen triggered by ER $\text{Ca}^{2+}$ depletion depends on cellular glutathione

Recent studies using fluorescent redox sensors targeted into the ER revealed that depletion of the organelle's  $\text{Ca}^{2+}$  store leads to a reductive shift in luminal redox balance in the time scale of minutes [13, 23, 24]. Either an irreversible inhibitor of SERCA (thapsigargin, TG) or the physiological  $\text{Ca}^{2+}$  mobilizing agents histamine [24] and cholecystokinin [23] rapidly transformed the ER luminal environment into a more reducing milieu. Although ER redox poise is known to control  $\text{Ca}^{2+}$  pumps and channels [37, 38], the relationship in the opposite direction, namely how  $\text{Ca}^{2+}$  can regulate the redox balance, has not been fully elucidated (for a recent review, see [39, 40]).

Given the close redox links between ER and mitochondria [41, 42], we initially evaluated the possibility of a mechanism that involves the mitochondria. However, neither mitochondrial superoxide production nor mitochondrial membrane potential or respiration was conspicuously affected by short-term (5–15 min) treatment with TG (Additional file 1: Fig. S1). What is more, ER stress over a longer time period leads to regulated protein reflux to the cytosol in budding yeast [43, 44]. Here, however, short-term ER  $\text{Ca}^{2+}$  depletion was not associated with the relocalization of the fluorescent redox sensor to the cytosol, as evidenced by co-staining immunofluorescence microscopy (Additional file 2: Fig. S2).

We further reasoned that such rapid change in redox poise could plausibly be explained by induced influx of reductants from the cytosol or efflux of oxidants to the cytosol. Because ER luminal redox poise strongly depends on the  $[\text{GSH}]^2:[\text{GSSG}]$  ratio, we first measured how the  $\text{Ca}^{2+}$  depletion-induced reductive shift was influenced by cellular glutathione levels. For monitoring ER redox poise, we used HEK293 cells stably expressing the specific  $[\text{GSH}]^2:[\text{GSSG}]$  sensor Grx1-roGFP1-iE<sub>ER</sub> [13]. Ratiometric measurements revealed that inhibition of SERCA by TG provoked a rapid reductive transition in Grx1-roGFP1-iE<sub>ER</sub> redox state in agreement with


previous results (Fig. 1a). However, when cellular GSH levels were depleted by overnight treatment with buthionine sulfoximine (BSO), the reductive transition upon addition of TG was abolished (Fig. 1b). BSO treatment resulted in a 75% drop of total glutathione concentration in HEK293 cells (Fig. 1c). We concluded each experiment by the consecutive addition of diamide and DTT to ensure the functionality of the probe. These results suggested that  $\text{Ca}^{2+}$  depletion-provoked reduction of the ER requires the cellular glutathione pool and residual glutathione in BSO-treated cells cannot mediate this process.

### Cytosolic Grx-roGFP2 is not detectably disturbed upon $\text{Ca}^{2+}$ release

In the ER lumen, the GSH/GSSG redox couple is shifted towards its oxidized form as a result of oxidative protein folding and the restricted permeability of GSSG through the ER membrane [16]. Accordingly, the quick reductive shift in response to TG could be caused by the flux of ER luminal GSSG to the cytosol, which would be expected to affect the cytosolic  $[\text{GSH}]^2:[\text{GSSG}]$  ratio. This possibility was tested by monitoring the redox state of the cytosolic  $[\text{GSH}]^2:[\text{GSSG}]$  sensor Grx1-roGFP2 [14] upon ER  $\text{Ca}^{2+}$  depletion. To prevent GSSG re-reduction by glutathione reductase (GR), Grx1-roGFP2-expressing HEK293 cells were pretreated with the GR-inhibitor carmustine (BCNU). We found that the redox poise of the cytosol was not measurably disturbed in response to TG (Fig. 2), suggesting that a mechanism other than GSSG efflux was responsible for the glutathione-dependent ER reduction.

### Permeabilization of the plasma membrane prevents the thapsigargin-induced reduction of the ER lumen

To further evaluate the possibilities of TG-induced ER import or export, we reasoned that global depletion of cytosolic components would affect the former possibility only. Digitonin selectively permeabilizes the plasma membrane due to its different lipid composition, but leaves intracellular membranes intact. Such treatment strongly dilutes the components of the cytosol and permits the examination of ER redox balance without cytosolic influence [18, 45]. The process of permeabilization was first visualized by monitoring the fluorescence decline in HeLa cells that were preloaded with BCECF-AM fluorescent dye (Fig. 3a). The persistence of the ER-localized fraction of the dye after 2 min of incubation with digitonin indicated the preserved integrity of the ER membrane (Fig. 3a). Using these optimized permeabilization conditions, HeLa cells were then transfected with HyPer-ER, permeabilized, and subjected to fluorescence ratio imaging. In this complex setup, we chose to use HyPer-ER rather than Grx1-roGFP1-iE<sub>ER</sub>



**Fig. 1**  $\text{Ca}^{2+}$  depletion-triggered ER reduction is sensitive to glutathione depletion by BSO. HEK293 cells were stably transfected with Grx1-roGFP1-iER constructs and subjected to ratiometric laser scanning microscopy on a temperature-controlled stage with  $\text{CO}_2$  control. Fluorescence ratio changes were monitored over time. Each trace corresponds to the data recorded from one cell; traces were obtained from two independent experiments. One micromolar TG were applied to untreated (**a**) or BSO-treated (**b**) cells as indicated by the arrow. At the end of each experiment, 500  $\mu\text{M}$  diamide (Dia) and 20 mM DTT were added to ensure the functionality of the probe. **c** Determination of total glutathione concentration by glutathione reductase assay as described in the “Materials and methods” section. One millimolar BSO treatment was performed overnight prior to experiment


because of its superior dynamic range [46]. It is important to emphasize that this non-specific redox sensor reliably monitors the process of TG-induced ER reduction [24] (Additional file 3: Fig. S3). Digitonin did not seem to influence the steady-state redox state of HyPer-ER but abolished the TG-induced luminal reduction (Fig. 3b). This observation suggested that the rapid hypo-oxidation is strongly dependent on a cytosolic component such as GSH and disqualified the speculated efflux of oxidizing molecules such as GSSG.

#### Thapsigargin increases glutathione levels in the ER lumen

We next analyzed possible changes in glutathione levels in the ER ( $[\text{GS}_{\text{tot}}]_{\text{ER}}$ ). To this end, we used a recently published method for calculation of  $[\text{GS}_{\text{tot}}]_{\text{ER}}$  [12] that combines the experimental values of  $[\text{GSH}]^2:[\text{GSSG}]$  and  $[\text{GSH}]:[\text{GSSG}]$  (Fig. 4a). Thus, we first determined  $[\text{GSH}]^2:[\text{GSSG}]$  in the ER by subjecting the Grx1-

roGFP1-iER-expressing HEK293 line to a quantitative plate-reader assay [25] before and after treatment with TG for 15 min. Consistent with the results above, ER  $[\text{GSH}]^2:[\text{GSSG}]$  rose from  $103 \pm 4$  to  $291 \pm 33$  mM upon treatment with TG (Fig. 4b). To determine ER  $[\text{GSH}]:[\text{GSSG}]$ , we transfected HEK293 cells with ER-targeted sGrx1p [12], which specifically equilibrates with  $[\text{GSH}]:[\text{GSSG}]$  [47]. As shown in Fig. 4c, the reduced sGrx1p:glutathionylated sGrx1p ratio ( $[-\text{SH}]:[-\text{SSG}]$ ) did not parallel the reductive shift of  $[\text{GSH}]^2:[\text{GSSG}]$  upon TG. In fact,  $[-\text{SH}]:[-\text{SSG}]$  rather decreased in response to treatment with TG.  $[-\text{SH}]:[-\text{SSG}]$  is proportional to  $[\text{GSH}]:[\text{GSSG}]$  [12] but, in the range below 0.1, can only be approximated by densitometry. We therefore only qualitatively concluded that ER  $[\text{GSH}]:[\text{GSSG}]$  remains constant or decreases in response to TG and that—according to the formula in Fig. 4a— $[\text{GS}_{\text{tot}}]_{\text{ER}}$  increases concomitantly to  $\text{Ca}^{2+}$  depletion-induced reduction of the ER.





Together with the above results using BSO, Grx1-roGFP2, and digitonin, these data strongly indicated that cytosolic GSH enters the ER lumen upon ER luminal  $\text{Ca}^{2+}$  depletion.

#### Chelation of cytosolic $\text{Ca}^{2+}$ does not inhibit glutathione transport

The depletion of ER  $\text{Ca}^{2+}$  is always accompanied by an increase in the cytosolic  $\text{Ca}^{2+}$  concentration. To resolve which side plays a role in the regulation of GSH transport, we buffered cytosolic  $\text{Ca}^{2+}$  with the chelator molecule BAPTA. Irrespective of hampered cytosolic  $\text{Ca}^{2+}$  fluxes, TG provoked the prompt reduction of the luminal GSH sensor, indicating that the decrease in luminal rather than the increase in cytosolic  $\text{Ca}^{2+}$  content triggers GSH transport (Fig. 5). These observations are consistent with the findings by Avezov et al. [23].

#### Cyclosporine A promotes GSH transport into the ER

Members of the cyclophilin family have been reported to be resident in the ER [48, 49]. They participate in the regulation of oxidative protein folding and ERAD [49, 50]. Moreover, their prototypic inhibitor cyclosporine A (CsA) causes an oxidative shift in cellular glutathione, presumably by increasing the oxidation state of the ER [49]. On this basis, we investigated if CsA treatment inhibits  $\text{Ca}^{2+}$  release-triggered GSH transport.


Unexpectedly, real-time monitoring of Grx1-roGFP1- $\text{iE}_{\text{ER}}$  revealed that CsA addition alone provoked the same immediate sensor reduction that was seen after TG addition (Fig. 6a). As for TG, the CsA-induced redox transition was sensitive to cellular GSH depletion (Fig. 6a) and short-term CsA treatment increased  $[\text{GS}_{\text{total}}]_{\text{ER}}$  (Fig. 4). Inhibition of both CsA- and TG-induced ER reduction by BSO strongly suggests a common, glutathione-centered mechanism.

The most thoroughly described mechanism of CsA action is the inhibition of the phosphatase activity of calcineurin, which prevents the activation of T lymphocytes [51]. CsA binds to the peptidyl-prolyl cis-trans isomerase cyclophilin A in the cytosol where the CsA-cyclophilin A complex mediates calcineurin inhibition [52]. To clarify, if calcineurin inhibition underlies ER reduction, we applied two mechanistically unrelated inhibitors of calcineurin, FK506 and cypermethrin [53]. Both calcineurin inhibitors failed to induce ER reduction, suggesting that the effect of CsA on ER GSH is independent of calcineurin (Fig. 6b, c). To test whether the GSH transporter might be directly gated by cyclophilins, we silenced the expression of cyclophilin A and the ER-resident cyclophilin B [48] and probed the sensor redox state after CsA addition. CsA-induced probe reduction was insensitive to the silencing of either cyclophilin (Fig. 6d), implying that CsA provokes GSH transport through another mechanism. Furthermore, although CsA is a well-known inhibitor of the mitochondrial permeability transition pore through blocking cyclophilin D [54], examination of its immediate effects on mitochondrial function showed only marginal changes (Additional file 1: Fig. S1).

#### GSH transport into the ER is not mediated by Sec61

Glutathione is present in every cellular compartment [55]. However, although there are several reports on GSH transport through the ER membrane [16–18], an ER GSH transporter has not yet been identified [56].

One possible candidate is the Sec61 translocon polypeptide channel, which allows the permeation of various small molecules through the ER membrane when not occupied by translocating polypeptide [34, 36]. Therefore, we examined whether opening the Sec61 channel would affect the ER luminal redox state. In agreement with earlier data [57], the application of puromycin, a translation inhibitor, which opens the Sec61 pore by clearing the nascent polypeptide, induced a comparable ER reduction as seen after TG addition (Fig. 7a). This reducing shift could be prevented by anisomycin (Fig. 7b), a known inhibitor of the unplugging action of puromycin [33]. Similar to TG, puromycin-induced ER reduction also depended on cellular glutathione levels, since BSO treatment or digitonin-mediated permeabilization of the plasma membrane abolished the



**Fig. 3** Permeabilization of the plasma membrane prevents thapsigargin-induced ER lumen reduction. **a** Sequential images of digitonin (25  $\mu$ g/ml)-treated HeLa cells loaded with BCECF-AM fluorescent dye. **b, c** Fluorescence ratio changes of HyPer-ER sensor 24 h after transfection in digitonin-permeabilized (red line) or intact (blue line) HeLa cells. Cells were pretreated with digitonin for 3 min and washed with intracellular medium as described in the “Materials and methods” section prior to the experiment. TG (200 nM, **b**) or puromycin (100  $\mu$ M, **c**) were applied at 3 min of imaging as indicated by the arrow. Experiments were terminated by addition of 0.5 mM DTT. Traces represent average intensity ratios acquired from 14 to 34 cells of 4 independent experiments

reducing shift (Figs. 7c and 3c), but had no apparent effects on mitochondrial function (Additional file 1: Fig. S1).


The translocon channel can also act as a  $\text{Ca}^{2+}$  leak channel [58]. Therefore, opening of the translocon can either trigger  $\text{Ca}^{2+}$  release and indirectly induce  $\text{Ca}^{2+}$ -sensitive GSH transport or directly facilitate the transport of GSH through the polypeptide channel itself. To distinguish between these two possibilities, we silenced Sec61 expression and examined the TG-induced redox change in Grx1-roGFP1-iE<sub>ER</sub>-expressing cells.  $\text{Ca}^{2+}$  depletion-dependent reduction was indistinguishable in Sec61-silenced and in non-silenced cells (Fig. 7d), suggesting that Sec61 was not directly involved in the transport of GSH.

We further examined whether  $\text{Ca}^{2+}$  depletion-induced reduction can be influenced by plugging the Sec61 translocon. Thus, cells were treated with anisomycin before TG addition. Since Sec61 is not the only possible  $\text{Ca}^{2+}$  leak channel in the ER membrane [34], we surmised that this treatment combination can further prove that Sec61

is dispensable for GSH transport. Indeed, sealing the Sec61 polypeptide channel with anisomycin did not prevent the TG-induced redox shift (Fig. 7e). We also applied puromycin before TG addition or in opposite order and observed no additive effect of the compounds in terms of hypo-oxidation of the ER lumen (Fig. 7f).

Kar2p, the yeast homolog of BiP, has recently been reported as a redox-dependent regulator of GSH influx into the ER through the Sec61 translocon [36]. Although our experiments in mammalian cells so far suggested that Sec61 was only indirectly involved in the inducible transport of GSH, we also checked for a possible regulation by BiP. However, neither the silencing of BiP nor its cleavage by subtilase toxin influenced the kinetics of ER reduction by CsA-induced GSH influx (Additional file 4: Fig. S4).


Together, this data argues that in mammalian cells, the Sec61 translocon does not participate in the  $\text{Ca}^{2+}$  depletion- or CsA-induced redox shift apart from serving as a  $\text{Ca}^{2+}$  leak channel in the presence of puromycin.



**Calreticulin is not required for the reduction of ER redox probes induced by Ca<sup>2+</sup> depletion**

Ca<sup>2+</sup> depletion hampers the mobility of the ER oxidoreductase PDI1A in the ER, which was explained by complex formation at low [Ca<sup>2+</sup>] between the Ca<sup>2+</sup>-binding chaperone CRT and PDI1A [28]. Moreover, the TG-

induced reductive shift, as measured by fluorescence lifetime of roGFPiE, appeared less prominent in CRT –/– mouse embryonic fibroblasts than in wild-type cells. It was concluded that the CRT-dependent decrease in mobility of PDI1A could be the mechanistic basis of Ca<sup>2+</sup> depletion-induced ER hypo-oxidation [28].






Figure 6.D


**Fig. 6** GSH transport can be triggered by cyclosporine A. **a** Real-time fluorescence ratio changes of Grx1-roGFP1-iER in response to 10 micromolar CsA in HEK293 cells stably expressing the sensor. Each trace corresponds to the data recorded from one cell; traces were obtained from two independent experiments. At the end of each experiment, 500  $\mu$ M diamide (Dia) and 20 mM DTT were added to ensure the functionality of the probe. Cells were left untreated or treated overnight with 1 mM BSO prior to the experiment. **b, c** Experiment performed as in **a**, but 50  $\mu$ M FK506 (**b**) or 10  $\mu$ M cypermethrin (**c**) was applied as marked instead of CsA. **d** HEK293 cells stably expressing Grx1-roGFP1-iER were transfected with control, cyclophilin A or B siRNA for 48 h before imaging; 10  $\mu$ M CsA were applied as indicated by the arrow. Knockdown efficiency was verified by qPCR

In light of our new findings that inducible ER hypo-oxidation depends on bulk import of GSH from the cytosol, we revisited the CRT hypothesis in our system. To this end, wild-type and CRT  $-/-$  mouse embryonic fibroblasts were transfected with Grx1-roGFP1-iER and analyzed by fluorescence video microscopy upon addition of TG, puromycin, or CsA. The fluorescence ratio curves of wild-type and CRT  $-/-$  cells equally responded at the time point of compound addition (Fig. 8), strongly indicating that CRT is dispensable for the induction of GSH-dependent ER reduction.

## Discussion

The maintenance of ER thiol-disulfide balance is of vital importance for the proper functioning of luminal processes, particularly the oxidative protein folding. Productive oxidative protein folding in the ER critically depends on the supply of disulfide reductants, which are required to resolve mispaired disulfide crosslinks in folding substrates [59, 60]. Currently, there is evidence for two cytosol-to-ER shuttling pathways for disulfide reductants: (i) a NADPH/thioredoxin reductase (TrxR)-






**Fig. 7** The Sec61 translocon polypeptide channel does not participate in glutathione transport. Effects of manipulating the translocon on fluorescence ratio changes of Grx1-roGFP1-iE<sub>ER</sub> in HEK293 cells stably expressing the sensor. Each trace corresponds to the data recorded from one cell. At the end of each experiment, 500  $\mu$ M diamide (Dia) and 20 mM DTT were added to ensure the functionality of the probe. **a** One hundred micromolar puromycin, **b** 200  $\mu$ M anisomycin followed by 100  $\mu$ M puromycin, **e** 200  $\mu$ M anisomycin followed by 1  $\mu$ M TG, and **f** 100  $\mu$ M puromycin followed by 1  $\mu$ M TG were applied as indicated by the arrow. **c** Cells were treated overnight with 1 mM BSO prior to experiment, and 100  $\mu$ M puromycin were applied as marked. **d** HEK293 cells stably expressing Grx1-roGFP1-iE<sub>ER</sub> were transfected with control or Sec61 siRNA for 48 h before imaging as above; 1  $\mu$ M TG were applied as indicated by the arrow. Knockdown efficiency was verified by Western blot (aSec61a, anti-Sec61a antibody; aBiP, anti-BiP antibody; aActin, anti-actin antibody)

dependent pathway [61] and (ii) a mechanism for GSH import into the ER [18, 62, 63]. In mammalian cells, both of these pathways are molecularly ill-defined, as is their presumable functional complementarity. It has recently been suggested that the TrxR-dependent pathway operates under non-stress conditions, whereas the GSH import pathway with its almost non-limited reducing capacity is mainly activated upon stress [59]. Indeed,

housekeeping protein reduction events during oxidative protein folding do not require ER luminal GSH [64], whereas the millimolar GSH pool in the ER [12] is instrumental for the non-catalyzed elimination of increased ER H<sub>2</sub>O<sub>2</sub> pools under stress [11].

Depletion of luminal Ca<sup>2+</sup>, either during normal cellular physiology or upon addition of pharmacological agents, provokes a rapid and reversible shift towards a more



**Fig. 8** Calreticulin is dispensable for ER reduction induced by  $\text{Ca}^{2+}$  depletion or Cyclosporin A. Wild-type and CRT  $-/-$  mouse embryonic fibroblasts were transfected with Grx1-roGFP1-iER<sub>+</sub>, and real-time fluorescent ratio changes were monitored. Reductive shift-provoking agents were applied as indicated. Each trace represents the data recorded from one cell; traces shown are representative of three independent experiments

reducing redox state of the luminal  $[\text{GSH}]^2:[\text{GSSG}]$  ratio [13, 23, 24].  $\text{Ca}^{2+}$  depletion-induced ER redox alterations are relevant in physiological conditions associated with  $\text{Ca}^{2+}$  signaling, such as the response of pancreatic cells to secretagogues and neuronal activity. Moreover, ER stress leads to a general decrease in luminal  $[\text{Ca}^{2+}]$  [41, 65], whereas the resulting rapid supply of thiols might help to dissolve stress-dependent protein aggregates and/or ER  $\text{H}_2\text{O}_2$  accumulation [11]. In principle, this redox shift could be due to (i) a transient decrease in the activity of luminal oxidases and/or oxidoreductases, (ii) the induction of a hypothetical luminal reductase, (iii) a transmembrane influx/efflux of reductants/oxidants, or (iv) a combination of these events. In this study, we present convincing evidence for the third possibility.

We found that  $\text{Ca}^{2+}$  depletion-induced luminal reduction requires the presence of cytosolic GSH: inhibition of GSH synthesis by BSO or the release of cytosolic GSH by selective permeabilization of the plasma membrane prevented the redox shift upon  $\text{Ca}^{2+}$  release. These observations suggested that GSH influx rather than GSSG efflux is responsible for the phenomenon. This interpretation was corroborated by the findings that the cytosolic redox state was not measurably changed after  $\text{Ca}^{2+}$  release from the ER and that  $[\text{GS}_{\text{tot}}]_{\text{ER}}$  was elevated rather than diminished in response to TG. Taken together, these results showed that GSH influx is the mechanism of  $\text{Ca}^{2+}$  depletion-induced luminal reduction. Of note, the steady-state sensor oxidation within the ER was not changed in BSO-treated cells. This is

consistent with previous findings that the redox state of PDI-family members does not change in response to BSO-mediated glutathione depletion [19]. The subcellular distribution of glutathione in BSO-treated cells is currently unclear and warrants further research.

$\text{Ca}^{2+}$  depletion also influences the mobility of PDIA1 via complex formation with the  $\text{Ca}^{2+}$ -binding chaperone CRT [28]. PDIA1 is the major ER oxidoreductase that shuttles newly generated disulfides to a variety of disulfide acceptors such as nascent protein folding substrates and GSH [8]. CRT-dependent immobilization of PDIA1 at low  $[\text{Ca}^{2+}]$  was proposed to explain the rapid ER reduction [28]. Consistently, the reductive shift of roGFP1iE that was induced by TG tended to be less prominent in CRT  $-/-$  mouse embryonic fibroblasts. This implies that in a  $\text{Ca}^{2+}$ -depleted environment, the binding of PDIA1 to CRT slows down the rate of oxidative protein folding, thus provoking ER hypo-oxidation. However, given the ER glutathione concentration of several millimolar [12] and its immediate response within approximately 3 min [13, 23, 24], an explanation that argues with lowered input of newly generated disulfides appears insufficient for kinetic reasons. Indeed, our experiments with Grx1-roGFP1-iE<sub>ER</sub> in wild-type and CRT  $-/-$  mouse embryonic fibroblasts showed equal responses to three reductive shift provoking agents, TG, puromycin, and CsA (Fig. 8). The discrepancy to the data from Avezov et al. could potentially be explained by the use of a glutathione-specific as opposed to a non-specific redox-sensing fluorescent protein reporter. The non-specific reporter used by Avezov et al. does not equilibrate with the glutathione redox couple [23] but may exhibit a certain selectivity to react with PDIA1 [66]. We conclude that CRT is dispensable for the rapid reduction of ER glutathione.

Glutathione biosynthesis resides exclusively in the cytosol [67], and glutathione transporters in intracellular membranes have not been identified at molecular level [55, 68]. However, functional studies revealed that GSH is able to cross the ER membrane, while permeation of GSSG is poor [16]. Since our work uncovers inducible GSH transport into the ER, we investigated the possible involvement of some candidate membrane proteins. The Sec61 translocon polypeptide channel has been reported to mediate the flux of some low molecular compounds beside proteins; however, the translocon opener puromycin did not enhance GSH transport significantly. A recent study postulated the translocon as an ER GSH transporter in yeast [36]. Indeed, the opening of the channel by puromycin reproduced the effect of TG on ER glutathione and the channel blocker anisomycin abolished the outcome, which also depended on the cytosolic GSH pool (Fig. 7a, b). However, silencing of Sec61 or plugging the channel with anisomycin did not

result in the inhibition of TG-induced ER reduction. These results suggest that the Sec61 translocon behaves as one of several types of  $\text{Ca}^{2+}$  leakage channels in the ER but does not directly participate in inducible GSH transport through the ER membrane in human cells.

We previously proposed a model, by which the passive ER influx of cytosol-derived GSH followed by its oxidation to membrane-impermeable GSSG will “lower  $[\text{GSH}]_{\text{ER}}$  and set up a driving force for further import of GSH from the cytosol. According to this model, the ER would constitute a trap for cellular glutathione, which is reminiscent of the mechanism of osmosis where an impermeable metabolite drives the diffusion of a permeable metabolite across a selectivity barrier such as a biological membrane.” [12]. The finding that GSH can enter the ER by facilitated diffusion through the Sec61 translocon in a yeast mutant [36] is congruent with this model, even though it should be acknowledged that the toxic ~ 10-fold increase in cytosolic GSH in this yeast mutant represents a rather non-physiological situation with regard to glutathione gradients at intracellular membranes. The present data now rather suggests an active as opposed to passive GSH import mechanism across the ER membrane. We are still hesitant, however, to conclusively dismiss the possibility of passive GSH import along a cytosol-to-ER  $[\text{GSH}]$  gradient that may be maintained opposite to the ER-to-cytosol  $[\text{GSSG}]_{\text{tot}}$  gradient reported earlier [12]. Such passive transport could be facilitated by a reversibly sealable, non-Sec61 permeation pore in the ER membrane.

We observed, much to our surprise, that CsA mimicked rather than inhibited the effect of TG on ER glutathione. Interestingly, cyclophilins, which are known targets of CsA, are involved in the regulation of the ER luminal milieu. On the one hand, overexpression of cytosolic cyclophilin A attenuates  $\text{Ca}^{2+}$  efflux from the ER, thereby inhibiting TG-induced apoptosis [69]. On the other hand, depletion of ER luminal cyclophilins results in ER hyper-oxidation with an elevated cellular GSSG:GSH ratio [49]. However, TG- and CsA-induced ER reduction was found to be independent of cyclophilins. The effect of CsA on ER glutathione was also independent of calcineurin, a prominent downstream target of the compound and a known modulator of ER  $\text{Ca}^{2+}$  channels [70]. CsA is also a prototypic inhibitor of glutathione- or glutathione conjugate-transporters of the ABC transporter superfamily, which operate in the plasma membrane [71]. However, our results showing a CsA-stimulated GSH influx into the ER speak against a possible involvement of ABC transporters.

Collectively, our data define a  $\text{Ca}^{2+}$ - and CsA-sensitive transport mechanism of GSH at the ER membrane. This transport does not involve the translocon polypeptide channel or CsA-sensitive ABC transporters. We also

excluded cyclophilins A and B, calcineurin, and CRT as regulatory components of GSH transport. Further studies are needed to explore this transport process in more detail.

## Conclusions

$\text{Ca}^{2+}$  mobilization from the ER results in influx of cytosolic GSH, which causes a redox shift towards more reducing conditions in the ER lumen. The mechanism may serve for the compensation of ER hyper-oxidation during excessive oxidative protein folding and/or ER stress. ER luminal redox-driven regulation of  $\text{Ca}^{2+}$  flux is well characterized and is known to involve inositol 1,4,5-trisphosphate receptors, ryanodine receptors, and sarco/endoplasmic reticulum  $\text{Ca}^{2+}$  transport ATPase [39, 40]. ER hyper-oxidation promotes  $\text{Ca}^{2+}$  release by the opening of ER  $\text{Ca}^{2+}$  channels and the inhibition of ER  $\text{Ca}^{2+}$  pumps. The present study unravels a homeostatic mechanism where  $\text{Ca}^{2+}$  depletion, in turn, can activate a GSH transporter, which will restore a proper ER redox environment (Fig. 9). This mechanism supports the feedback regulation of oxidative protein folding and contributes to the robustness of ER luminal redox balance.

## Materials and methods

### Generation of HEK293 cells stably expressing Grx1-roGFP1-iE<sub>ER</sub>

HEK293 cells were transfected with Grx1-roGFP1-iE<sub>ER</sub>/pcDNA3.1 [13] using Metafectene PRO (Biontex) and stably expressing clones selected by addition of 1 mg/ml

G418 (Sigma). Homogeneous expression of clones was checked by fluorescence microscopy at the excitation wavelength of 405 nm. Clone D5 was selected for further experiments.


### Cell culture and transient transfections

HeLa and HEK293 cells were cultured in Dulbecco's modified Eagle's medium (DMEM) (Invitrogen) containing 4.5 g/l glucose supplemented with 10% fetal bovine serum, 100 U/ml penicillin, and 100 mg/ml streptomycin at 37 °C in 5%  $\text{CO}_2$ . For cells stably expressing Grx1-roGFP1-iE<sub>ER</sub>, G418 (1 mg/ml) was added to the growth medium as a selection antibiotic.

Transient transfections with cytosolic Grx-roGFP2 [14] or HyPer-ER constructs [24] were performed with Lipofectamine (Thermo Fisher) reagent according to manufacturer's instructions; cells were analyzed 48 h after transfection.

For silencing Sec61, HEK293 cells stably expressing Grx1-roGFP1-iE<sub>ER</sub> were transfected using Lipofectamine RNAiMax reagent based on the manufacturer's protocols, using final concentration of 100 nm of siRNA. Negative control and *SEC61A1* siRNA were previously published [58]. Successful knockdown was confirmed by Western blot analysis using anti-Sec61α primary antibody [58].

Silencing cyclophilin A and B siRNA was delivered by Lipofectamine RNAiMax (Thermo Fisher Scientific) according to manufacturer's recommendation; 85 pM siRNA and 2.5 μl Lipofectamine reagent were used per 50,000 cells. The target sequence of the mock siRNA was 5'-UGGUUUACAUGUUUCUGA-3', of the cyclophilin A



**Fig. 9** Schematic representation of feedback loops that connect ER  $\text{Ca}^{2+}$  loading, GSH influx, and oxidative protein folding. Hyper-oxidizing conditions in the ER (orange box) due to peak oxidative protein folding leads to  $\text{Ca}^{2+}$  depletion via opening of IP3R calcium channels and inhibition of SERCA pumps.  $\text{Ca}^{2+}$  depletion can in turn activate a GSH transporter (yellow box), which will restore the proper steady-state ER redox environment (green box). Conversely, hyper-reducing conditions in the ER (blue box) lower GSH influx via increased  $[\text{Ca}^{2+}]_{\text{ER}}$ , thereby rescuing steady-state ER redox and commensurate oxidative protein folding. These feedback mechanisms regulate the pace of oxidative protein folding and contribute to the robustness of ER luminal redox balance



siRNA was 5'-CUGGAUUGCAGAGUUAAGU-3', and of the cyclophilin B siRNA 5'-CAAAAACAGUGGAUAAUUU-3' (Microsynth, Switzerland).

#### Quantitative PCR and gene expression analysis

To assess gene expression, total RNA was extracted using TRI reagent (Sigma). Subsequently, cDNA was produced by reverse transcription with Maloney murine leukemia virus reverse transcriptase (Promega). Quantitative PCR (qPCR) analysis was performed using the KAPA SYBR Fast kit (Sigma) on a Rotor Gene Real-Time Cycler (Corbett Research). Normalization of the data relative to the endogenous control gene glyceraldehyde-3-phosphate dehydrogenase (GAPDH) was done according to the  $2^{-\Delta\Delta C_t}$  method for relative quantification.

Primers:

##### 1. Human cyclophilin A

- (a) FW: CAT CTG CAC TGC CAA GAC TGA
- (b) Rev: TGC AAT CCA GCT AGG CAT G

##### 2. Human cyclophilin B

- (a) FW: GGT GAT CTT TGG TCT CTT CGG
- (b) Rev: TAG ATG CTC TTT CCT CCT GTG

##### 3. GAPDH

- (a) FW: TGA TGA CAT CAA GAA GGT GGT GAA
- (b) Rev: TCC TTG GAG GCC ATG TGG GCC AT

#### Cultivation and transfection of mouse embryonic fibroblasts

The control and CRT  $-/-$  MEFs were kindly provided by Maurizio Molinari (Bellinzona, Switzerland) with the kind permission of Marek Michalak (Edmonton, Canada) [72]. The cells were cultured in  $\alpha$ MEM containing 10% FBS and 100 U/ml penicillin and 0.1 mg/ml streptomycin under standard culture conditions (37 °C, 5% CO<sub>2</sub>). DNA transfection was performed using Xfect (Takara) according to manufacturer's instructions; cells were analyzed 48 h after transfection.

#### Live-cell imaging of Grx1-roGFP1-iE<sub>ER</sub>

Live-cell imaging was performed on an Olympus Fluoview 1000 (experiments for Fig. 8: Olympus Fluoview 3000) laser scanning confocal microscope equipped with a  $\times 60$  (experiments for Fig. 8,  $\times 40$ ) oil immersion objective (NA 1.40), a 405-nm laser diode, a 440 nm (experiments for Fig. 8, 445 nm) laser diode, and a 488-nm argon gas laser. The 405- and 440/445-nm laser lines were used as excitation wavelengths; the emission window was set to 500–600 nm. Images were acquired in sequential frame mode, separating the two channels. Grx1-roGFP1-iE<sub>ER</sub>-expressing cells were grown on glass bottom dishes (Mattek); for ratiometric analysis, cells were washed twice with DMEM without phenol red and transferred to a heated chamber (37 °C) with CO<sub>2</sub> control. Reagents were added in 1 ml phenol red-free DMEM in the required concentration. For Ca<sup>2+</sup> chelation experiments, cells were pretreated for 30 min with 50  $\mu$ M BAPTA-AM. At the end of each experiment, 500  $\mu$ M diamide and 20 mM DTT were added. Images were taken every 30 s for a period of 30 min and analyzed with the ImageJ software. One region of interest (ROI) per cell was chosen, which remained immobile for the duration of image acquisition, and 405/440 ratios were determined from emission intensities in background subtracted ROIs.

#### Live-cell imaging of HyPer-ER

HeLa cells were analyzed 48 h after HyPer-ER transfection by fluorescent excitation ratiometry. Fluorescence-intensity measurements were performed on an inverted microscope (Axio Observer, Zeiss) equipped with a  $40\times 1.4$  oil-immersion objective (Fluar, Zeiss) and a Cascade II camera (Photometrics, Tucson, AZ). Excitation wavelengths were set by a random-access monochromator connected to a xenon arc lamp (DeltaRAM, Photon Technology International, Birmingham, NJ). For ratiometric measurements of HyPer-ER, excitation wavelengths of 490 and 420 nm were sequentially applied combined with a 505-nm dichroic filter and a 525/36-nm emission filter set. Cells grown on 10 cm coverslips were washed with HEPES-buffered solution containing 145 mM NaCl, 5 mM KCl, 1 mM MgCl<sub>2</sub>, 0.8 mM CaCl<sub>2</sub>, 10 mM HEPES, 5 mM glucose, and pH 7.4 and placed into a heated chamber at 37 °C. Reagents were added in 10 $\times$  concentration in 0.1 ml of prewarmed buffer after removing 0.1 ml of medium. At the end of each experiment, 20 mM DTT was added to check sensor sensitivity. Images were acquired every 10 s for a period of 30 min and analyzed by the MetaFluor (Molecular Devices, Woburn, MA) software. Oxidation state of HyPer-ER was calculated by 490/420-nm fluorescence excitation ratio of HyPer-ER after background fluorescence subtraction.

For plasma membrane permeabilization, cells were treated with digitonin (25  $\mu$ g/ml) for 3 min prior to

experiment and washed with intracellular (IC) medium containing 113.5 mM KCL, 5 mM NaHCO<sub>3</sub>, 4 mM MgCl<sub>2</sub>, 40 nM CaCl<sub>2</sub>, 5 mM K-EGTA, 20 mM HEPES, 4 mM ATP, and 5.6 mM D-glucose. Experiments were performed after signal stabilization.

#### Measurement of [GSH]<sup>2</sup>:[GSSG] and [GSH]:[GSSG] in the ER

To estimate [GS<sub>tot</sub>]<sub>ER</sub>, we used the procedure published in Montero et al. [12]. The degree of oxidation (OxD) of Grx1-roGFP1-iE<sub>ER</sub> was quantitatively determined in cells stably expressing Grx1-roGFP1-iE<sub>ER</sub> in 96-well plates (Falcon) in complete medium without phenol red. One day after seeding, cells were treated with 1 μM thapsigargin and 10 μM CsA, or left untreated for 15 min. The completely oxidized and reduced conditions were achieved by adding 500 μM diamide or 10 mM DTT to each pretreatment respectively before excitation spectrum analysis. Fluorescent intensities were measured on 520 nm from the bottom on Spectramax Gemini EM (Molecular Device) in a 350–500-nm range. OxD values and OxD-derived [GSH]<sup>2</sup>:[GSSG] values were calculated as published before [25].

The glutathionylation status of sGrx1p<sub>ER</sub> was analyzed in transiently transfected HEK293 cells by densitometric analysis of [<sup>35</sup>S]-methionine metabolically labeled, alkylated, and immunoprecipitated protein as described previously [12]. Cells were left untreated or treated with 1 μM thapsigargin or 10 μM CsA for 15 min ahead of analysis.

#### Supplementary information

**Supplementary information** accompanies this paper at <https://doi.org/10.1186/s12915-020-0749-y>.

**Additional file 1: Figure S1.** Short-term thapsigargin, cyclosporine A and puromycin treatment has only minor effects on mitochondrial superoxide production, the mitochondrial membrane potential, and mitochondrial respiration. (A) Mitochondrial superoxide production was measured in HEK cells seeded at a density of 30000 cells per well to a 96 well plate the day before measurement. Cells were loaded with 5 μM MitoSOX red and Hoechst 33342 (both Life Technologies) for 10 minutes. The cells were treated with either 0.1% DMSO, 30% ethanol (positive control), 10 μM cyclosporine A (CsA), 1 μM thapsigargin, or 100 μM puromycin, and mitochondrial superoxide production was measured for the indicated times on a Cellomics ArrayScan VTI HCS Reader (Thermo Scientific). (B) Mitochondrial membrane potential was detected in HEK cells seeded at a density of 30000 cells per well to a 96 well plate the day before measurement. For positive control, cells were treated with 230 nM valinomycin (Sigma) for six hours. The cells were treated with either 0.1% DMSO, 10 μM cyclosporine A, 1 μM thapsigargin or 100 μM puromycin for 15 to 60 minutes in complete medium. The cells were stained for 30 minutes with Hoechst 33342 and MitoTracker Red CMXRos (both Life Technologies) and analyzed on a Cellomics ArrayScan VTI HCS Reader counting at least 1000 cells per well. Values were normalized to 15 minutes DMSO. (C) Left panel: Real-time measurements of oxygen consumption rate (OCR), reflecting mitochondrial respiration were performed on a Seahorse XF96 Analyzer (Agilent Technologies, USA) based on previous description (Nagy et al., *Biochim Biophys Acta Bioenerg* 2018, 1859, 201–214). Where indicated by arrows, cells were treated with metabolic inhibitors/modulators (oligomycin 2 μM, FCCP 100 nM, and antimycin A + rotenone 1 μM

each). TG, puromycin, and CsA were applied 5 min ahead of the recording at the following concentrations: 1 μM, 100 μM, and 10 μM, respectively. Right panel: Basal respiration rate was detected before oligomycin addition; ATP-linked respiration was calculated as a difference between the values before oligomycin and before carbonyl cyanide-4-(trifluoromethoxy)phenylhydrazone (FCCP) addition; maximal respiratory capacity is the difference between values before and after rotenone + antimycin A addition; spare capacity difference between values before oligomycin and before rotenone + antimycin A addition; non-mitochondrial respiration is shown by the last value after rotenone + antimycin A addition. Data were obtained from two independent experiments (mean ± s.d.). (PPTX 258 kb)

**Additional file 2: Figure S2.** Grx1-roGFP1-iE<sub>ER</sub> is not released from the ER upon treatment of cells with thapsigargin, cyclosporine A, or puromycin for 30 min. HEK293 cells stably expressing HA-tagged Grx1-roGFP1-iE<sub>ER</sub> were treated with 0.1 %DMSO, 10 μM cyclosporine A (CsA), 1 μM thapsigargin (TG) or 100 μM puromycin (Puro) for 30 min. Cells were fixed, stained with Hoechst 33342, incubated with anti-Calnexin and anti-HA antibodies followed by green- and red-fluorescent secondary antibodies, and analysed on an epifluorescence microscope. Overlay images including increased magnification frames of selected insets are shown for each treatment in colour. Size bar, 10 μm. (PPTX 2915 kb)

**Additional file 3: Figure S3.** BSO treatment prevents Hyper-ER reduction upon addition of thapsigargin. Fluorescence ratio changes of Hyper-ER sensor 24 hours after transfection in untreated or BSO treated HeLa cells. (PPTX 5979 kb)

**Additional file 4: Figure S4.** BiP does not influence the Ca<sup>2+</sup> depletion-dependent reductive shift in the ER. Effect of manipulation of BiP on fluorescence ratio changes of Grx1-roGFP1-iE<sub>ER</sub> in HEK293 cells stably expressing the sensor. BiP levels were diminished by the addition of subtilaseAB (SubAB) toxin or by silencing (BiP kd). As control, an inactive subtilaseA<sub>A272</sub>B mutant (SubAB mut) and a non-silencing control siRNA (control kd) were also used. The effect of subtilaseAB treatment (A) or BiP silencing (B) was checked by immunoblotting. (C) Identical curves of reductive shift were observed upon CsA addition in all cases; SubAB treatment for 60 min, siRNA transfection for 48 h. (PPTX 1031 kb)

#### Abbreviations

BSO: Buthionine sulfoximine; CsA: Cyclosporine A; ER: Endoplasmic reticulum; TG: Thapsigargin

#### Acknowledgements

This study is dedicated to the memory of Gábor Bánhegyi (1957–2019). We would like to thank Mrs. Valéria Mile for her skilful technical assistance, Marek Michalak and Maurizio Molinari for the kind gift of control and CRT –/– MEFs, and James and Adrienne Paton, University of Adelaide, Australia, for the generous gift of subtilaseAB toxin.

#### Authors' contributions

BL contributed to the experimental design and analysis, performance of experiments, data acquisition and analysis, and writing of the original draft of the manuscript. JB and CAH contributed to the performance of experiments, and data acquisition and analysis. MZ, GK, and DVK performed a set of experiments. AO and RZ contributed to the experimental design and analysis and revision of the manuscript. CAH and GB contributed to the experimental design of the study, data analyses, and writing of the manuscript. All authors read and approved the final manuscript.

#### Funding

This work was supported by grants from the National Research, Development and Innovation Office, Hungary (111899 to BL, 119955 to MG, 112696 and 124813 to GB). CA-H was a recipient of an Ambizione fellowship by the Swiss National Science Foundation.

#### Availability of data and materials

The datasets used and/or analyzed during the current study were deposited on Zenodo [73]. Reagents specific to this study are available on request.

**Ethics approval and consent to participate**

Not applicable.

**Consent for publication**

Not applicable.

**Competing interests**

The authors declare that they have no competing interests.

**Author details**

<sup>1</sup>Department of Medical Chemistry, Molecular Biology and Pathobiochemistry, Semmelweis University, Budapest, Hungary. <sup>2</sup>Division of Molecular and Systems Toxicology, Department of Pharmaceutical Sciences, University of Basel, Klingelbergstrasse 50, 4056 Basel, Switzerland. <sup>3</sup>Department of Physiology, Faculty of Medicine, Semmelweis University, Budapest, Hungary. <sup>4</sup>"Momentum" Peroxidase Enzyme Research Group of the Semmelweis University and the Hungarian Academy of Sciences, Budapest, Hungary. <sup>5</sup>Medical Biochemistry and Molecular Biology, Saarland University, 66421 Homburg, Germany. <sup>6</sup>University Medical Library, University of Basel, Spiegelgasse 5, 4051 Basel, Switzerland.

Received: 17 December 2019 Accepted: 11 February 2020

Published online: 26 February 2020

**References**

- Braakman I, Bulleid NJ. Protein folding and modification in the mammalian endoplasmic reticulum. *Annu Rev Biochem.* 2011;80:71–99.
- Csala M, Kereszturi E, Mandl J, Banhegyi G. The endoplasmic reticulum as the extracellular space inside the cell: role in protein folding and glycosylation. *Antioxid Redox Signal.* 2012;16(10):1100–8.
- Margittai E, Sitia R. Oxidative protein folding in the secretory pathway and redox signaling across compartments and cells. *Traffic.* 2011;12(1):1–8.
- Hetz C, Papa FR. The unfolded protein response and cell fate control. *Mol Cell.* 2018;69(2):169–81.
- Camello C, Lomax R, Petersen OH, Tepikin AV. Calcium leak from intracellular stores—the enigma of calcium signalling. *Cell Calcium.* 2002;32(5–6):355–61.
- Anelli T, Bergamelli L, Margittai E, Rimessi A, Fagioli C, Malgaroli A, Pinton P, Ripamonti M, Rizzuto R, Sitia R. Ero1alpha regulates Ca(2+) fluxes at the endoplasmic reticulum-mitochondria interface (MAM). *Antioxid Redox Signal.* 2012;16(10):1077–87.
- Prins D, Michalak M. Organellar calcium buffers. *Cold Spring Harb Perspect Biol.* 2011;3(3):a004069.
- Bulleid NJ, Ellgaard L. Multiple ways to make disulfides. *Trends Biochem Sci.* 2011;36(9):485–92.
- Appenzeller-Herzog C. Glutathione- and non-glutathione-based oxidant control in the endoplasmic reticulum. *J Cell Sci.* 2011;124(Pt 6):847–55.
- Lappi AK, Ruddock LW. Reexamination of the role of interplay between glutathione and protein disulfide isomerase. *J Mol Biol.* 2011;409(2):238–49.
- Melo EP, Lopes C, Gollwitzer P, Lortz S, Lenzen S, Mehmeti I, Kaminski CF, Ron D, Avezov E. TriPer, an optical probe tuned to the endoplasmic reticulum tracks changes in luminal H<sub>2</sub>O<sub>2</sub>. *BMC Biol.* 2017;15(1):24.
- Montero D, Tachibana C, Rahr Winther J, Appenzeller-Herzog C. Intracellular glutathione pools are heterogeneously concentrated. *Redox Biol.* 2013;1:508–13.
- Birk J, Meyer M, Aller I, Hansen HG, Odermatt A, Dick TP, Meyer AJ, Appenzeller-Herzog C. Endoplasmic reticulum: reduced and oxidized glutathione revisited. *J Cell Sci.* 2013;126(Pt 7):1604–17.
- Gutscher M, Pauleau AL, Marty L, Brach T, Wabnitz GH, Samstag Y, Meyer AJ, Dick TP. Real-time imaging of the intracellular glutathione redox potential. *Nat Methods.* 2008;5(6):553–9.
- Hwang C, Sinskey AJ, Lodish HF. Oxidized redox state of glutathione in the endoplasmic reticulum. *Science.* 1992;257(5076):1496–502.
- Banhegyi G, Lusini L, Puskas F, Rossi R, Fulceri R, Braun L, Mile V, di Simplicio P, Mandl J, Benedetti A. Preferential transport of glutathione versus glutathione disulfide in rat liver microsomal vesicles. *J Biol Chem.* 1999;274(18):12213–6.
- Jessop CE, Bulleid NJ. Glutathione directly reduces an oxidoreductase in the endoplasmic reticulum of mammalian cells. *J Biol Chem.* 2004;279(53):55341–7.
- Molteni SN, Fassio A, Ciriolo MR, Filomeni G, Pasqualetto E, Fagioli C, Sitia R. Glutathione limits Ero1-dependent oxidation in the endoplasmic reticulum. *J Biol Chem.* 2004;279(31):32667–73.
- Appenzeller-Herzog C, Riemer J, Zito E, Chin KT, Ron D, Spiess M, Ellgaard L. Disulfide production by Ero1alpha-PDI relay is rapid and effectively regulated. *EMBO J.* 2010;29(19):3318–29.
- Margittai E, Enyedi B, Csala M, Geiszt M, Banhegyi G. Composition of the redox environment of the endoplasmic reticulum and sources of hydrogen peroxide. *Free Radic Biol Med.* 2015;83:331–40.
- Hudson DA, Gannon SA, Thorpe C. Oxidative protein folding: from thiol-disulfide exchange reactions to the redox poise of the endoplasmic reticulum. *Free Radic Biol Med.* 2015;80:171–82.
- Banhegyi G, Benedetti A, Csala M, Mandl J. Stress on redox. *FEBS Lett.* 2007;581(19):3634–40.
- Avezov E, Cross BC, Kaminski Schierle GS, Winters M, Harding HP, Melo EP, Kaminski CF, Ron D. Lifetime imaging of a fluorescent protein sensor reveals surprising stability of ER thiol redox. *J Cell Biol.* 2013;201(2):337–49.
- Enyedi B, Varnai P, Geiszt M. Redox state of the endoplasmic reticulum is controlled by Ero1L-alpha and intraluminal calcium. *Antioxid Redox Signal.* 2010;13(6):721–9.
- Birk J, Ramming T, Odermatt A, Appenzeller-Herzog C. Green fluorescent protein-based monitoring of endoplasmic reticulum redox poise. *Front Genet.* 2013;4:108.
- Ermakova YG, Bilan DS, Matlashov ME, Mishina NM, Markvicheva KN, Subach OM, Subach FV, Bogeski I, Hoth M, Enikolopov G, et al. Red fluorescent genetically encoded indicator for intracellular hydrogen peroxide. *Nat Commun.* 2014;5:5222.
- Mehmeti I, Lortz S, Lenzen S. The H<sub>2</sub>O<sub>2</sub>-sensitive HyPer protein targeted to the endoplasmic reticulum as a mirror of the oxidizing thiol-disulfide milieu. *Free Radic Biol Med.* 2012;53(7):1451–8.
- Avezov E, Konno T, Zyryanova A, Chen W, Laine R, Crespiello-Casado A, Melo E, Ushioda R, Nagata K, Kaminski CF, et al. Retarded PDI diffusion and a reductive shift in poise of the calcium depleted endoplasmic reticulum. *BMC Biol.* 2015;13(1):2.
- Vishnu N, Jadoon Khan M, Karsten F, Groschner LN, Waldeck-Weiermair M, Rost R, Hallstrom S, Imamura H, Graier WF, Malli R. ATP increases within the lumen of the endoplasmic reticulum upon intracellular Ca<sup>2+</sup>-release. *Mol Biol Cell.* 2014;25(3):368–79.
- Klein MC, Zimmermann K, Schorr S, Landini M, Klemens PAW, Altensell J, Jung M, Krause E, Nguyen D, Helms V, et al. AXER is an ATP/ADP exchanger in the membrane of the endoplasmic reticulum. *Nat Commun.* 2018;9(1):3489.
- Yong J, Bischof H, Burgstaller S, Siirin M, Murphy A, Malli R, Kaufman RJ. Mitochondria supply ATP to the ER through a mechanism antagonized by cytosolic Ca(2+). *eLife.* 2019;8:e49682.
- Csala M, Marcolongo P, Lizak B, Senesi S, Margittai E, Fulceri R, Magyar JE, Benedetti A, Banhegyi G. Transport and transporters in the endoplasmic reticulum. *Biochim Biophys Acta.* 2007;1768(6):1325–41.
- Lizak B, Czeglé I, Csala M, Benedetti A, Mandl J, Banhegyi G. Translocon pores in the endoplasmic reticulum are permeable to small anions. *Am J Physiol Cell Physiol.* 2006;291(3):C511–7.
- Lizak B, Csala M, Benedetti A, Banhegyi G. The translocon and the non-specific transport of small molecules in the endoplasmic reticulum (review). *Mol Membr Biol.* 2008;25(2):95–101.
- Schauble N, Lang S, Jung M, Cappel S, Schorr S, Ulucan O, Linxweiler J, Dudek J, Blum R, Helms V, et al. BiP-mediated closing of the Sec61 channel limits Ca<sup>2+</sup> leakage from the ER. *EMBO J.* 2012;31(15):3282–96.
- Posner AJ, Igbaria A, Darch MA, Miled S, Outten CE, Winther JR, Palais G, D'Auteaux B, Delaunay-Moisano A, Toledano MB. Endoplasmic reticulum transport of glutathione by Sec61 is regulated by Ero1 and Bip. *Mol Cell.* 2017;67(6):962–73 e965.
- Higo T, Hattori M, Nakamura T, Natsume T, Michikawa T, Mikoshiba K. Subtype-specific and ER luminal environment-dependent regulation of inositol 1,4,5-trisphosphate receptor type 1 by ERp44. *Cell.* 2005;120(1):85–98.
- Li Y, Camacho P. Ca<sup>2+</sup>-dependent redox modulation of SERCA 2b by ERp57. *J Cell Biol.* 2004;164(1):35–46.
- Appenzeller-Herzog C, Simmen T. ER-luminal thiol/selenol-mediated regulation of Ca<sup>2+</sup> signalling. *Biochem Soc Trans.* 2016;44(2):452–9.
- Chernorudskiy AL, Zito E. Regulation of calcium homeostasis by ER redox: a close-up of the ER/mitochondria connection. *J Mol Biol.* 2017;429(5):620–32.

41. Eletto D, Chevet E, Argon Y, Appenzeller-Herzog C. Redox controls UPR to control redox. *J Cell Sci.* 2014;127(Pt 17):3649–58.
42. Fan Y, Simmen T. Mechanistic connections between endoplasmic reticulum (er) redox control and mitochondrial metabolism. *Cells.* 2019;8(9):1071.
43. Lajoie P, Snapp EL: Size-dependent secretory protein reflux into the cytosol in association with acute endoplasmic reticulum stress. *bioRxiv* 2019:573428.
44. Igbaria A, Merksamer PJ, Trusina A, Tilahun F, Johnson JR, Brandman O, Krogan NJ, Weissman JS, Papa FR. Chaperone-mediated reflux of secretory proteins to the cytosol during endoplasmic reticulum stress. *Proc Natl Acad Sci.* 2019;116(23):11291–8.
45. Wilson R, Allen AJ, Oliver J, Brookman JL, High S, Bulleid NJ. The translocation, folding, assembly and redox-dependent degradation of secretory and membrane proteins in semi-permeabilized mammalian cells. *Biochem J.* 1995;307(Pt 3):679–87.
46. Belousov VV, Fradkov AF, Lukyanov KA, Staroverov DB, Shakhbazov KS, Tersikh AV, Lukyanov S. Genetically encoded fluorescent indicator for intracellular hydrogen peroxide. *Nat Methods.* 2006;3(4):281–6.
47. Iversen R, Andersen PA, Jensen KS, Winther JR, Sigurskjold BW. Thiol-disulfide exchange between glutaredoxin and glutathione. *Biochemistry.* 2010;49(4):810–20.
48. Price ER, Zydowsky LD, Jin MJ, Baker CH, McKeon FD, Walsh CT. Human cyclophilin B: a second cyclophilin gene encodes a peptidyl-prolyl isomerase with a signal sequence. *Proc Natl Acad Sci U S A.* 1991;88(5):1903–7.
49. Stocki P, Chapman DC, Beach LA, Williams DB. Depletion of cyclophilins B and C leads to dysregulation of endoplasmic reticulum redox homeostasis. *J Biol Chem.* 2014;289(33):23086–96.
50. Bernasconi R, Solda T, Galli C, Pertel T, Luban J, Molinari M. Cyclosporine A-sensitive, cyclophilin B-dependent endoplasmic reticulum-associated degradation. *PLoS One.* 2010;5(9):e13008.
51. Fruman DA, Klee CB, Bierer BE, Burakoff SJ. Calcineurin phosphatase activity in T lymphocytes is inhibited by FK 506 and cyclosporin A. *Proc Natl Acad Sci U S A.* 1992;89(9):3686–90.
52. Handschumacher RE, Harding MW, Rice J, Drugge RJ, Speicher DW. Cyclophilin: a specific cytosolic binding protein for cyclosporin A. *Science.* 1984;226(4674):544–7.
53. Enan E, Matsumura F. Specific inhibition of calcineurin by type II synthetic pyrethroid insecticides. *Biochem Pharmacol.* 1992;43(8):1777–84.
54. Griffiths EJ, Halestrap AP. Further evidence that cyclosporin A protects mitochondria from calcium overload by inhibiting a matrix peptidyl-prolyl cis-trans isomerase. Implications for the immunosuppressive and toxic effects of cyclosporin. *Biochem J.* 1991;274(Pt 2):611–4.
55. Banerjee R. Redox outside the box: linking extracellular redox remodeling with intracellular redox metabolism. *J Biol Chem.* 2012;287(7):4397–402.
56. Bachhawat AK, Thakur A, Kaur J, Zulkifli M. Glutathione transporters. *Biochim Biophys Acta.* 2013;1830(5):3154–64.
57. van Lith M, Tiwari S, Pediani J, Milligan G, Bulleid NJ. Real-time monitoring of redox changes in the mammalian endoplasmic reticulum. *J Cell Sci.* 2011; 124(Pt 14):2349–56.
58. Lang S, Schauble N, Cavalie A, Zimmermann R. Live cell calcium imaging combined with siRNA mediated gene silencing identifies Ca(2+)-leak channels in the ER membrane and their regulatory mechanisms. *J Visualized Exp.* 2011;53:e2730.
59. Delaunay-Moisan A, Ponsero A, Toledano MB. Reexamining the function of glutathione in oxidative protein folding and secretion. *Antioxid Redox Signal.* 2017;27(15):1178–99.
60. Ellgaard L, Sevier CS, Bulleid NJ. How are proteins reduced in the endoplasmic reticulum? *Trends Biochem Sci.* 2018;43(1):32–43.
61. Poet GJ, Oka OB, van Lith M, Cao Z, Robinson PJ, Pringle MA, Arner ES, Bulleid NJ. Cytosolic thioredoxin reductase 1 is required for correct disulfide formation in the ER. *EMBO J.* 2017;36(5):693–702.
62. Chakravarthy S, Bulleid NJ. Glutathione is required to regulate the formation of native disulfide bonds within proteins entering the secretory pathway. *J Biol Chem.* 2004;279(38):39872–9.
63. Cuozzo JW, Kaiser CA. Competition between glutathione and protein thiols for disulfide-bond formation. *Nat Cell Biol.* 1999;1(3):130–5.
64. Tsunoda S, Avezov E, Zyryanova A, Konno T, Mendes-Silva L, Pinho Melo E, Harding HP, Ron D. Intact protein folding in the glutathione-depleted endoplasmic reticulum implicates alternative protein thiol reductants. *eLife.* 2014;3:e03421.
65. Kiviluoto S, Vervliet T, Ivanova H, Decuyper JP, De Smedt H, Missiaen L, Bultynck G, Parys JB. Regulation of inositol 1,4,5-trisphosphate receptors during endoplasmic reticulum stress. *Biochim Biophys Acta.* 2013;1833(7): 1612–24.
66. Konno T, Pinho Melo E, Lopes C, Mehmeti I, Lenzen S, Ron D, Avezov E. ERO1-independent production of H<sub>2</sub>O<sub>2</sub> within the endoplasmic reticulum fuels Prdx4-mediated oxidative protein folding. *J Cell Biol.* 2015;211(2):253–9.
67. Lu SC. Glutathione synthesis. *Biochim Biophys Acta.* 2013;1830(5):3143–53.
68. Oestreicher J, Morgan B. Glutathione: subcellular distribution and membrane transport (1). *Biochem Cell Biol.* 2019;97(3):270–89.
69. Kim J, Choi TG, Ding Y, Kim Y, Ha KS, Lee KH, Kang I, Ha J, Kaufman RJ, Lee J, et al. Overexpressed cyclophilin B suppresses apoptosis associated with ROS and Ca<sup>2+</sup> homeostasis after ER stress. *J Cell Sci.* 2008;121(Pt 21):3636–48.
70. Cameron AM, Steiner JP, Roskams AJ, Ali SM, Ronnett GV, Snyder SH. Calcineurin associated with the inositol 1,4,5-trisphosphate receptor-FKBP12 complex modulates Ca<sup>2+</sup> flux. *Cell.* 1995;83(3):463–72.
71. Ballatori N, Hammond CL, Cunningham JB, Krance SM, Marchan R. Molecular mechanisms of reduced glutathione transport: role of the MRP/CFTR/ABCC and OATP/SLC21A families of membrane proteins. *Toxicol Appl Pharmacol.* 2005;204(3):238–55.
72. Mesaeli N, Nakamura K, Zvaritch E, Dickie P, Dziak E, Krause KH, Opas M, MacLennan DH, Michalak M. Calreticulin is essential for cardiac development. *J Cell Biol.* 1999;144(5):857–68.
73. Lizák B, Birk J, Zana M, Kosztyi G, Kratschmar DV, Odermatt A, et al. Ca<sup>2+</sup> mobilization-dependent reduction of the endoplasmic reticulum lumen is due to influx of cytosolic glutathione. 2020, Supporting Datasets. <https://zenodo.org/record/3648401>

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

